

The Authors declare no conflicts of interest.

References

- 1) Daniels G. *Human Blood Groups* [M], 2nd edn. Oxford, UK: Blackwell Science Press; 2002. p. 165.
- 2) Chang JG, Wang JC, Yang TY, et al. Human RhD is caused by a deletion of 1,013 bp between introns 8 and 9 including exon 9 of *RHD* gene. *Blood* 1998; **92**: 2602-4.
- 3) Shao CP, Maas JH, Su YQ, et al. Molecular background of RhD-positive, D-negative, D(e) and weak D phenotypes in Chinese. *Vox Sang* 2002; **83**: 156-61.
- 4) Lin IL, Shih MC, Hsieh MH, et al. Molecular basis of weak D in Taiwanese. *Ann Hematol* 2003; **82**: 617-20.
- 5) Xu Q, Zhang J, Wang Q, et al. *RHD* gene polymorphism among RhD-negative Han Chinese. *Chin Med J (Engl)* 2003; **116**: 1539-43.
- 6) Shao CP, Xiong W. A new hybrid *RHD* positive, D antigen negative allele. *Transfus Med* 2004; **14**: 185-6.
- 7) Xu Q, Grootkerk-Tax MG, Maaskant-Van Wijk PA, et al. Systemic analysis and zygosity determination of the *RHD* gene in a D-negative Chinese Han population reveals a novel D-negative *RHD* gene. *Vox Sang* 2005; **88**: 35-40.
- 8) Wu JJ, Hong XZ, Xu XG, et al. Molecular basis of partial D phenotypes in Chinese [in Chinese]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi* 2006; **14**: 587-91.
- 9) Yan L, Wu J, Zhu F, et al. Molecular basis of D variants in Chinese persons. *Transfusion* 2007; **47**: 471-7.
- 10) Sun G, Jing H, Xiong W, et al. A weak D type 12 found in the Chinese. *Chin J Blood Transfus* 2006; **19**: 14-7.
- 11) Ye LY, Guo ZH, Li Q, Zhu ZY. Molecular and family analyses revealed two novel *RHD* alleles in a survey of a Chinese RhD-negative population. *Vox Sang* 2007; **92**: 242-6.
- 12) Ye LY, Guo ZH, Zhu ZY. Homo sapiens Rhesus blood group D antigen (*RHD*) gene, *RHD*-weak D allele, exon 1 and partial cds (EU335051), Homo sapiens Rhesus blood group D antigen (*RHD*) gene, *RHD*-weak D allele, exon 9 and partial cds (EF103573) and Homo sapiens Rhesus blood group D antigen (*RHD*) gene, *RHD*(A414V) allele, exon 10 and partial cds (EU604751). Available at: <http://www.ncbi.nlm.nih.gov/genbank>. Accessed on 10/07/2013.
- 13) Ye L, Yue D, Wo D, et al. Molecular bases of unexpressed *RHD* alleles in Chinese D- persons. *Transfusion* 2009; **49**: 1655-60.
- 14) Li Q, Hou L, Guo ZH, et al. Molecular basis of the *RHD* gene in blood donors with DEL phenotypes in Shanghai. *Vox Sang* 2009; **97**: 139-46.
- 15) Xu H, Ye SH, Shao CP, et al. A new *RHD*-positive, D antigen negative allele in Chinese. *Vox Sang* 2010; **98**: 576.
- 16) The Ministry of Public Health of China. Guidelines of technical operation for transfusion. 2nd edn. Tianjin, China: Tianjin Science and Technology Press; 1998. p. 64.
- 17) Shao CP, Xiong W. A new PCR method for direct determination of *RHD* zygosity. *Natl Med J China* 2004; **84**: 736-9.
- 18) Legler TJ, Maas JH, Kohler M, et al. *RHD* sequencing: a new tool for decision making on transfusion therapy and provision of Rh prophylaxis. *Transfus Med* 2001; **11**: 383-8.
- 19) Wagner FF, Frohmajer A, Flegel WA. *RHD* positive haplotypes in D negative Europeans. *BMC Genet* 2001; **2**: 10.
- 20) Sun GD, Yi ZZ, Zhang YP, et al. *RHD* gene polymorphism of 118 Han population IAT D-negative individuals [in Chinese]. *Chin J Blood Transfus* 2007; **20**: 292-4.
- 21) Zhuo CS, Zhuo X, Guo YJ, et al. *RHD* gene polymorphism of RhD negative individuals in population of Fujian Province [In Chinese]. *J Exp Hematol* 2008; **16**: 435-8.
- 22) Cruz BR, Chiba AK, Moritz E, et al. *RHD* alleles in Brazilian blood donors with weak D or D-negative phenotypes. *Transfus Med* 2012; **22**: 84-9.
- 23) Omi T, Takahashi J, Tsudo N, et al. The genomic organization of the partial D category DVa: the presence of a new partial D associated with the DVa phenotype. *Biochem Biophys Res Commun* 1999; **254**: 786-94.
- 24) Qing Wei. Random survey for RH allele polymorphism among 50 native Tibetans. Dissertation for the attainment of the Doctor Degree of Medicine (Dr. med.) at the Faculty of Medicine, University of Ulm. 2006. Available at: http://vts.uni-ulm.de/query/longview.meta.asp?document_id=5775. Accessed on 10/07/2013.
- 25) Shao CP, Wang BY, Ye SH, et al. Childbearing-aged women and anti-D positive blood recipients prevent DEL transfusion from allosensitization and ineffective treatment in East Asia. *J Zhejiang Univ Sci B* 2012; **13**: 913-8.
- 26) Kim KH, Kim KE, Woo KS, et al. Primary anti-D immunization by DEL red blood cells. *Korean J Lab Med* 2009; **29**: 361-5.
- 27) Yasuda H, Ohto H, Sakuma S, et al. Secondary anti-D immunization by Del red blood cells. *Transfusion* 2005; **45**: 1581-4.
- 28) Shao CP. Transfusion of RhD-positive blood in "Asia type" DEL recipients. *N Engl J Med* 2010; **362**: 472-3.
- 29) Shao CP, Xu H, Xu Q, et al. Antenatal Rh prophylaxis is unnecessary for "Asia type" DEL women. *Transfusion Clin Bio* 2010; **17**: 260-4.

Arrived: 15 April 2013 - Revision accepted: 11 July 2013

Correspondence: Chao-Peng Shao
Department of Transfusion
The Second People's Hospital of Shenzhen
Shungang West Road
Futian, Shenzhen 518035, China
e-mail: cpshaos@hotmail.com