

- 27) Pelc-Klopotowska M, Guz K, Walaszcyk A, et al. Weak D type two patient with allo anti - D. *Vox Sang* 2010; **99**: 367.
- 28) Vege S, Fong C, Lomas-Francis C, et al. Weak D type 2 and production of anti-D. *Transfusion* 2011; **51**: 41A.
- 29) Pham BN, Roussel M, Peyrard T, et al. Anti-D investigations in individuals expressing weak D type 1 or weak D type 2: allo- or autoantibodies? *Transfusion* 2011; **51**: 2679-85.
- 30) Dogic V, Bingulac-Popovic J, Babic I, et al. Distribution of weak D types in the Croatian population. *Transfus Med* 2011; **21**: 278-9.
- 31) Christiansen M, Samuelsen B, Christiansen L, et al. Correlation between serology and genetics of weak D types in Denmark. *Transfusion* 2008; **48**: 187-93.
- 32) Flegel WA. How I manage donors and patients with a weak D phenotype. *Curr Opin Hematol* 2006; **13**: 476-83.
- 33) Ouchari M, Chakroun T, Abdelkefi S, et al. Anti-D auto-immunization in a patient with weak D type 4.0. *Transfus Clin Biol* 2014; **21**: 43-6.
- 34) Daniels G. *Human Blood Groups*. 2nd ed. Oxford: Blackwell Science; 2002. p. 195-274.
- 35) Ceppellini R, Dunn LC, Turry M. An interaction between alleles at the Rh locus in man which, weakens the reactivity of the Rh0 factor (Du). *Proc Natl Acad Sci USA* 1955; **41**: 283.
- 36) Touinssi M, Chapel-Fernandes S, Granier T, et al. Molecular analysis of inactive and active RHD alleles in native Congolese cohorts. *Transfusion* 2009; **49**: 1353-60.
- 37) Ouchari M, Jemni-Yaacoub S, Chakroun T, et al. *RHD* alleles in the Tunisian population. *Asian J Transfus Sci* 2013; **7**: 119-24.

Arrived: 18 March 2014 - Revision accepted: 29 May 2014

Correspondence: Saloua Jemni Yacoub
Centre Régionale de Transfusion Sanguine
Hôpital Universitaire Farhat Hached - Sousse
Rue Ibn Al Jazzar
Sousse, Tunisia
e-mail: saloua.jemni@rs.tn

© SIMTI Servizi Srl